

AQUATIC ANIMAL DISEASE REPORT - 2025

Country/territory: NEW ZEALAND

Item	Disease status/occurrence code a/c/												Level of diagnosis	Epidemiological comment numbers
DISEASES PREVALENT IN THE REGION	Month													
FINFISH DISEASES	January	February	March	April	May	June	July	August	September	October	November	December		
WOAH-listed diseases														
1. Infection with epizootic haematopoietic necrosis virus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
2. Infection with infectious haematopoietic necrosis virus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
3. Infection with spring viremia of carp virus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
4. Infection with viral haemorrhagic septicaemia virus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
5. Infection with <i>Aphanomyces invadans</i> (EUS)	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
6. Infection with Megalocytivirus pagrus 1 (formerly red sea bream iridovirus)	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
7. Infection with koi herpesvirus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
8. Infection with tilapia lake virus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
Non WOAH-listed diseases														
9. Grouper iridoviral disease	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
10. Viral encephalopathy and retinopathy	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
11. Enteric septicaemia of catfish	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
12. Carp Edema Virus Disease	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
MOLLUSC DISEASES														
WOAH-listed diseases														
1. Infection with <i>Bonamia exitiosa</i>	- (2024)	+	- (2025)	- (2025)	- (2025)	- (2025)	- (2025)	- (2025)	- (2025)	- (2025)	- (2025)	- (2025)		1
2. Infection with <i>Perkinsus olseni</i>	- (2024)	- (2024)	- (2024)	- (2024)	- (2024)	- (2024)	- (2024)	+?	- (2025)	- (2025)	- (2025)	- (2025)		2
3. Infection with abalone herpesvirus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
4. Infection with <i>Xenohaliotis californiensis</i>	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
5. Infection with <i>Bonamia ostreae</i>	- (2024)	-(2024)	- (2024)	-(2024)	- (2024)	+?	- (2025)	- (2025)	- (2025)	- (2025)	+?	+?		3
Non WOAH-listed diseases														
6. Infection with <i>Marteilioides chungmuensis</i>	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
7. Acute viral necrosis (in scallops)	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
CRUSTACEAN DISEASES														
WOAH-listed diseases														
1. Infection with Taura syndrome virus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
2. Infection with white spot syndrome virus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
3. Infection with yellow head virus genotype 1	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
4. Infection with infectious hypodermal and haematopoietic necrosis virus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
5. Infection with infectious myonecrosis virus	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
6. Infection with <i>Macrobrachium rosenbergii</i> nodavirus (White Tail disease)	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
7. Infection with <i>Hepatobacter penaei</i> (Necrotising hepatopancreatitis)	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
8. Acute hepatopancreatic necrosis disease (AHPND)	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
9. Infection with <i>Aphanomyces astaci</i> (Crayfish plague)	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
10. Infection with decapod iridescent virus 1 (DIV1)	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
Non WOAH-listed diseases														
11. Hepatopnacreatic Microsporidiosis caused by <i>Enterocytozoon hepatopenaei</i> (HPM-EHP)	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
12. Viral covert mortality disease (VCMD) of shrimps	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
13. <i>Spiroplasma eriocheiris</i> infection	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		
AMPHIBIAN DISEASES														
WOAH-listed diseases														
1. Infection with <i>Ranavirus</i> species	0000	0000	0000		0000	0000	0000	0000	0000	0000	0000	0000		
2. Infection with <i>Batrachochytrium dendrobatidis</i>	- (2022)	- (2022)	- (2022)	- (2022)	- (2022)	+?	- (2025)	- (2025)	- (2025)	- (2025)	- (2025)	- (2025)		4
3. Infection with <i>Batrachochytrium salamandrivorans</i>	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000	0000		

Prepared by:

Name: Rissa Williams and Giulia Raponi

Position: Senior Surveillance Advisor and Incursion Investigator

ANY OTHER DISEASES OF IMPORTANCE													
1													
2													

DISEASES PRESUMED EXOTIC TO THE REGION^b**LISTED BY THE WOA^H**

Finfish: Infection with HPR-deleted or HPR0 salmon anaemia virus; Infection with salmon pancreas disease virus;
Infection with *Gyrodactylus salaris* .

Molluscs: Infection with *Marteilia refringens* ; *Perkinsus marinus* .

NOT LISTED BY THE WOA^H

Finfish: Channel catfish virus disease

a/ Please use the following occurrence code:

<u>Occurrence code and symbol</u>	<u>Definition</u>	<u>Occurrence code and symbol</u>	<u>Definition</u>
Disease present +	The disease is present with clinical signs in the whole country (in domestic species or wildlife)	Disease absent -	The disease was absent in the country during the reporting period (in domestic species or wildlife).
Disease limited to one or more zones +()	The disease is present with clinical signs, and limited to one or more zones/compartments (in domestic species or wildlife)	Never reported 0000	The disease has "never been reported" (historically absent) for the whole country in domestic species and wildlife.
Infection/infestation +?	Confirmed infestation or infection using diagnostic tests, but no clinical signs observed (in domestic species or wildlife)	No information ***	No information is available regarding the presence or the absence of this disease during the reporting period (in domestic species or wildlife).
Infection/infestation limited to one or more zones +?()	Confirmed infestation or infection using diagnostic tests, but no clinical signs observed and limited to one or more zones/compartments (in domestic species or wildlife)		
Disease suspected ?	The presence of the disease was suspected but not confirmed (in domestic species or wildlife)		
Disease suspected but not confirmed and limited to one or more zones ?()	The presence of the disease was suspected but not confirmed and limited to one or more zones/compartments (in domestic species or wildlife)		

b/ If there is any changes on historical data, please highlight in RED

1. Epidemiological comments:

(Comments should include: 1) Origin of the disease or pathogen (history of the disease); 2) Species affected; 3) Disease characteristics (unusual clinical signs or lesions); 4) Pathogen (isolated/sero-typed); 5) Mortality rate (high/low; decreasing/increasing); 6) Death toll (economic loss, etc); 7) Size of infected areas or names of infected areas; 8) Preventive/control measures taken; 9) Samples sent to national or international laboratories for confirmation (indicate the names of laboratories); 10) Published paper (articles in journals/website, etc). and 11) Unknown diseases: describe details as much as possible.)

Comment No.	
1	<i>Bonamia exitiosa</i> occurs in wild flat oyster (<i>Ostrea chilensis</i>) beds in Foveaux Strait, Southland where it is highly prevalent and associated with mortalities in mid to late summer. It occurs intermittently around the South Island and the North Island of New Zealand. <i>Bonamia exitiosa</i> has been previously detected in flat oysters from Hauraki Gulf (Auckland region), Tauranga (Bay of Plenty region), the Marlborough Sounds and Wellington Harbour (Southern end of the North Island). Annual monitoring of the presence of <i>B. exitiosa</i> infection is undertaken in the flat oyster population in the Foveaux Strait, and in February 2025, 20.7% of the flat oysters tested positive for <i>B. exitiosa</i> , compared with 18.1% in 2024. The survey also estimated higher mortality associated with higher prevalence of <i>B. exitiosa</i> across the survey area.

2	1. Reported from the Nelson region via passive surveillance surveillance; 2. Species affected – wild trough clams (<i>Cyclomactra ovata</i>) 3. Clinical signs – n/a 4. Pathogen – <i>Perkinsus olseni</i> 5. Mortality rate – n/a 6. Economic loss – n/a 7. Geographic extent – Rabbit Island, Nelson 8. Containment measures – n/a; 9. Laboratory confirmation – real-time <i>Perkinsus</i> genus PCR, conventional <i>Perkinsus</i> genus PCR and nucleotide sequencing, <i>P. olseni</i> real-time PCR (National Animal Health Laboratory); 10. Publications – https://www.sciquest.org.nz/browse/publications/article/176644 11) Unknown diseases: describe details as much as possible.): n/a <i>Perkinsus olseni</i> was first detected in New Zealand in 1999, in wild wedge shells (<i>Macomona lilian</i> a). It was then found in wild populations of New Zealand cockles (<i>Austrovenus stutchburyi</i>), ark shells (<i>Barbatia novaezelandiae</i>) and pipi (<i>Paphies australis</i>) in 2000-2001. In July 2013, <i>P. olseni</i> was detected for the first time in farmed black foot pāua (<i>Haliotis iris</i>), an abalone species endemic to New Zealand. Further detections were made in wild <i>H. iris</i> populations in 2014. These mollusc species occur widely around the coast of New Zealand, but to date <i>P. olseni</i> has only been detected in these species from the Auckland region northwards. <i>Perkinsus olseni</i> was found for the first time on the South Island in New Zealand green lipped mussels (<i>Perna canaliculus</i>) in a land based aquaculture facility in September 2014, and then in wild New Zealand scallops (<i>Pecten novaezelandiae</i>) in November 2014. Both of these findings were in the Marlborough region, and were incidental and not associated with mortality events. In November 2017, passive surveillance detected <i>P. olseni</i> from New Zealand scallops in two sites within Kaipara harbour, Auckland region, and again was thought to be incidental and not associated with significant pathology in scallops. In August 2018, there was another incidental finding of <i>P. olseni</i> in farmed green lipped mussels (<i>Perna canaliculus</i>) in the Coromandel region (North Island), that was not associated with mortalities. In October 2019, <i>P. olseni</i> was detected in <i>P. canaliculus</i> in a land based aquaculture facility in Nelson that was experiencing low level mortalities. It remains unknown if <i>P. olseni</i> was related to the mortalities in <i>P. canaliculus</i> in this case. In January - March 2021, <i>P. olseni</i> was reported from a targeted survey of farmed greenlipped mussels in the Coromandel (Waikato region) and in Pelorus Sound (Marlborough region) and in wild greenlipped mussels in the Nelson region. Further detections of <i>P. olseni</i> were made during routine surveillance of broodstock green lipped mussels in a land based facility in November - December 2021 and March 2024. These detections were not associated with unusual mortalities. In August 2025, <i>P. olseni</i> was detected through molecular methods in 2/10 wild trough clams (<i>Cyclomactra ovata</i>) that were tested following a mass mortality event in the Nelson region. <i>Perkinsus olseni</i> was not, however, detected via histology any in any of the clams submitted. It was concluded that <i>P. olseni</i> was an incidental finding, and not a contributing factor to the mortality event, which was most likely triggered by preceding adverse environmental conditions.
3	1. Reported from Big Glory Bay via targeted surveillance; 2. Species affected – wild flat oysters (<i>Ostrea chilensis</i>) 3. Clinical signs – n/a 4. Pathogen – <i>Bonamia ostreae</i> 5. Mortality rate – n/a 6. Economic loss – n/a 7. Geographic extent – Big Glory Bay, Stewart Island (Southland) 8. Containment measures – n/a; 9. Laboratory confirmation – – ddPCR (National Institute Water and Atmospheric Research), qPCR and nucleotide sequencing (National Animal Health Laboratory); 10. Publications – n/a 11) Unknown diseases: describe details as much as possible.): n/a <i>Bonamia ostreae</i> was detected for the first time in New Zealand flat oysters (<i>Ostrea chilensis</i>) in January 2015. It was found in two regions in the northern part of the South Island: on one land-based aquaculture facility in the Nelson region, and on two marine farms in the Marlborough region. Since that time, movement controls have been in place to regulate the movement of susceptible shellfish from the northern regions of the South Island and active surveillance has been conducted for the purposes of early detection of spread. In 2016, <i>B. ostreae</i> was detected in both farmed and wild flat oysters within the Marlborough region (the same region as initially reported), and was associated with pathology and mortality in the farmed population. In May 2017 surveillance detected <i>B. ostreae</i> in marine flat oyster farms in Big Glory Bay, Stewart Island (situated in the Southland region, at the southern end of the South Island). No clinical signs or elevated mortality was observed in association with <i>B. ostreae</i> in farmed flat oysters in Big Glory Bay. Following this detection, movement controls to manage risk movements from Stewart Island were issued, and depopulation of all flat oyster farms within areas where <i>B. ostreae</i> had been detected commenced. Depopulation of farms in Big Glory Bay commenced on the 19 June 2017 and was completed in September 2017. Depopulation of farms in Marlborough Sounds commenced on the 11 July 2017 and was completed in December 2017. Targeted surveillance of wild flat oysters collected from Big Glory Bay in Targeted surveillance of wild flat oysters from Big Glory Bay in June 2025 detected <i>B. ostreae</i> in 169/220 animals collected in June 2025 and in 77/80 animals collected in November-December 2025. These detections were not associated with mortality.
4	The first isolation of <i>Batrachochytrium dendrobatidis</i> was made in 1999 in New Zealand. Since then the fungus has been detected both on the North and South Islands in both native and introduced frog species.

5	
2. New aquatic animal health regulations introduced within past six months (with effective date):	