



World Organisation
for Animal Health



中国动物疫病预防控制中心
China Animal Disease Control Center



MALAYSIA

Member's updates on the main diseases in domestic swine population—— **Nipah**

NUR AIMI SYARINA BINTI PAUZI
RESEARCH OFFICER
VETERINARY RESEARCH INSTITUTE (VRI)

DR. LEE CHIAN HOEY
VETERINARY OFFICER
DEPARTMENT OF VETERINARY SERVICES





The Sixth WOAHP Regional Training on Swine Disease
Control in Asia and the Pacific
Beijing, People's Republic of China
13-16 October 2026



Member's profile

2

Domestic Pig Population in Malaysia, 2017-2026





RISK OF INCURSION AND SPREAD OF MAJOR SWINE DISEASES IN MALAYSIA



HEALTHY
ENVIRONMENT



HEALTHY
ANIMALS



HEALTHY
PEOPLE



A SAFER
MALAYSIA



AFRICAN SWINE FEVER (ASFV)

HIGH RISK

Multiple entry and spread pathways, including animal movement, contaminated materials, trade and wild-pig interfaces.



Animal movement
and transport



Trade and
contaminated pork products



Contaminated materials
(equipment, vehicle, personnel)



Wild-pig interface
(domestic and wild pigs)



NIPAH VIRUS (NiV)

INTERMEDIATE RISK

Persistent fruit-bat reservoir and bat-pig interface, but risk is reduced by strengthened biosecurity, surveillance and movement control.



Fruit-bat reservoir
(Pteropus spp.)



Bat-pig interface
(near fruit trees/orchards)



Farm biosecurity
(open housing increases risk)



Reduced risk with
surveillance and
movement control



OTHER MAJOR SWINE DISEASES (e.g. CSFV, PRRSV, PCV, JE, SIV)

INTERMEDIATE RISK

Risk driven mainly by **animal movement, trade, farm biosecurity and regional disease pressure.**



Animal movement
and transport



Trade and
international/regional
disease pressure



Farm biosecurity
and management



Multiple circulating
swine pathogens
in the region



Malaysia – Nipah Virus Encephalitis

Disease significance

Nipah virus infection is a **zoonotic disease** and a **WOAH-listed disease**.



Malaysia experienced the first recognised Nipah virus outbreak in **1998–1999**.

The outbreak involved pig farms and human encephalitis cases.



Natural reservoir

Fruit bats (*Pteropus* spp.) are the natural reservoir hosts of Nipah virus.



Amplifying host

During the Malaysian outbreak, domestic pigs acted as an **amplifying host** for Nipah virus.



Nipah virus


Zoonotic transmission

The virus can be transmitted between animals and humans, causing severe disease in people (encephalitis).



Current status

Malaysia maintains **historical freedom from infection with Nipah virus**.





Disease situations - Historical outbreak

5



The Diagnostic Trap



Initial misdiagnosis of JE resulted in inappropriate vector control measures and a catastrophic 4-month delay in targeted intervention.



The Direct Link

Amplification occurs rapidly in high-density swine populations before jumping to humans.

NIPAH OUTBREAK: IMPACT ON PIG POPULATION IN PENINSULAR MALAYSIA



Pig movements carried the virus from Perak to other parts of Peninsular Malaysia (such as Bukit Pelanduk in Negeri Sembilan and Sepang, Selangor) and even to abattoir workers in Singapore.

FROM OUTBREAK CONTROL TO MAINTAINING FREEDOM

Current Nipah virus situation in Malaysia



1999

Outbreak controlled



2001

Self-declaration of
freedom published
by WOAH

2026

Freedom maintained
through continuous vigilance



1

NO ACTIVE DISEASE



Malaysia remains **free**
from Nipah virus.

2

**CONTINUOUS
SURVEILLANCE**



Annual surveillance
in **domestic pigs**
and **wild pigs**.

3

**PREPAREDNESS &
PREVENTION**



Early detection



Biosecurity



Import controls



Rapid reporting



2025 SURVEILLANCE

>2,000
domestic pig
samples

48
wild pig
samples



ALL NEGATIVE



Disease freedom is not the end of surveillance – **it is maintained by surveillance.**

Sources: DVS Malaysia, 2026; Malaysia's WOAH self-declaration, 2025



Response to disease incursion

7

INTEGRATED STRATEGIES TO PREVENT, DETECT AND RESPOND TO NIPAH VIRUS

1 IMPORT CONTROL

- Risk-based import requirements
- Quarantine and health certification




2 SURVEILLANCE

- Active and passive surveillance
- Laboratory testing (RT-PCR, ELISA, SNT)



3 REPORTING

- Mandatory and timely notification
- Timely reporting through ADIC for national coordination



4 OUTBREAK MANAGEMENT

- Farm status classification and zoning
- Movement control and follow-up testing



5 BIOSECURITY

- Prevent pig-bat and farm-to-farm transmission
- Control farm access and strengthen farm hygiene



6 OCCUPATIONAL SAFETY

- Appropriate PPE and safe animal handling
- Protect veterinarians, farm and abattoir workers



7 DISINFECTION

- Routine cleaning and disinfection
- Safe disposal of waste and carcasses



8 ADVICE

- Technical guidance for farmers and veterinarians
- Evidence-based biosecurity and risk reduction advice



9 PUBLIC AWARENESS

- Educate farmers, communities and stakeholders
- Promote early recognition and reporting



ONE HEALTH APPROACH



ANIMAL HEALTH



HUMAN HEALTH



ENVIRONMENT & WILDLIFE



Laboratory capacity

8



LABORATORY CAPACITY

VRI's Diagnostic Capability for Major Swine Diseases



HEALTHY
ENVIRONMENT



HEALTHY
ANIMALS



HEALTHY
PEOPLE



A SAFER
MALAYSIA



MOLECULAR DETECTION

- Conventional RT-PCR
- Nested / Semi-nested RT-PCR
- Real-time RT-PCR (qRT-PCR)
- Multiplex RT-PCR




GENETIC CHARACTERISATION

- Sanger sequencing (partial gene sequencing)
- Whole-genome sequencing (WGS)
- Phylogenetic and molecular epidemiology analysis




SEROLOGY (ANTIBODY DETECTION)

- Enzyme-Linked Immunosorbent Assay (ELISA)
- Serum Neutralisation Test (SNT)



CELL CULTURE (VIRUS ISOLATION)

- Virus isolation in suitable cell lines
- Observation of cytopathic effect (CPE)
- Confirmation by RT-PCR/sequencing



MAJOR SWINE DISEASES COVERED



**Nipah virus
(NiV)**



**African swine fever
virus (ASFV)**



**Classical swine fever
virus (CSFV)**



**Porcine reproductive
and respiratory syndrome
virus (PRRSV)**



**Porcine circovirus
(PCV)**



**Other swine viruses
(e.g. JE, SIV)**





Collaboration with other stakeholders

10



The Sixth WOA Regional Training on Swine Disease Control in Asia and the Pacific

13 – 16 October, 2026 Beijing, China

Challenge and possible solutions

11

CHALLENGES TO CONTROL NIPAH IN MALAYSIA AND MALAYSIA'S ACTIONS TO OVERCOME THOSE CHALLENGES



KEY CHALLENGES

Factors contributing to the challenges

1 Movement of infected or at-risk pigs

- Some farmers moved or sold pigs to other farms or markets when animals showed signs of illness to avoid financial losses from death or government culling.
- Difficult to trace animal movements.



2 High-density and intensive pig farming

- Large number of pigs kept in close proximity in certain areas increased the risk of rapid virus amplification and spread.
- Fast within-farm transmission.
- Greater impact during outbreaks.



3 Socio-economic impact and farmer compliance

- Culling of infected and in-contact pigs caused major financial losses, leading to reluctance to report, non-compliance with control measures and illegal movement of animals.



MALAYSIA'S ACTIONS

Strategies and measures implemented to overcome the challenges

1 Control of animal movement and traceability



- Movement control within and between states/areas.
- Quarantine and licensing for animal transport.
- Trace-back and trace-forward investigations.
- Enforcement by DVS with support from other agencies (e.g. police, local authorities).



2 Strengthened surveillance and early detection



- Active surveillance in pig farms, wildlife (bats) and environment.
- Enhanced diagnostic capacity (including BSL-3 for Nipah at VRI).
- Network of DVS laboratories across Malaysia.
- Improved disease reporting and information sharing.



3 Farmer awareness and support



- Continuous awareness campaigns on disease recognition.
- Guidance on reporting and not moving sick animals.
- Compensation for culled animals to reduce financial burden.
- Training on biosecurity and good farming practices.



FUTURE LONG-TERM DIRECTION FOR THE MALAYSIAN PIG INDUSTRY

FROM
CONVENTIONAL FARMING

TO
MODERN, BIOSECURE & SUSTAINABLE PIG FARMING



MODERN PIG FARMING (MPF) AND PIG FARMING AREAS (PFA)

Transform farms • strengthen biosecurity • reduce disease risk



Enhanced
biosecurity



Improved
farm management



Better waste
management



Sustainable
pig industry

DVS STRATEGIC PLAN 2022 – 2030 | KEY PRIORITIES

1



DISEASE FREEDOM

Work towards freedom
from ASF

2



SURVEILLANCE

Active surveillance +
MADIC reporting

3



PREPAREDNESS

Simulation exercises +
risk assessment

4



DIAGNOSTICS

ISO 17025 +
WOAH-aligned testing

5



TRACEABILITY & BORDER CONTROL

Traceability + import risk
& border controls



World Organisation
for Animal Health

Thank you!

Contact information



The Sixth WOA Regional Training on Swine Disease
Control in Asia and the Pacific
Beijing, People's Republic of China
13-16 October 2026