



African Swine Fever (ASF) Outbreak Investigation in Neverland, September-December, 2021

Vietnamese team:

Long Thanh Pham

Chuong Dinh Vo

Dzung Kim Nguyen



Objectives & Expected outcomes

1. To identify likely **source of ASFv introduction** to the regions
2. To find out the **transmission routes** between households/villages/regions
3. To identify possible **risk factors** associated with the ASF outbreak, and **types of farms** and regions are at **higher/lower risk** of ASF
4. To estimate approximate **economic loss** may be due to ASF
5. To propose recommendations for control and prevention of the outbreaks

Target group and case definition

- A cross-sectional survey was conducted between September - December 2021 in **3 administrative regions** (assumption) with 10 ASF outbreaks ongoing
- **Target group**
 - Backyard households
 - Commercial farms
 - Wild pigs
- **Case definition**
 - **Animal level:**
 - Acute haemorrhagic fever ($>40^{\circ}\text{C}$), reddening of the skin (white pigs) – tips of ears, tail, distal extremities, ventral aspects of chest and abdomen, severe lethargy, diarrhoea (sometime bloody), vomiting and eye discharge, sudden death, high mortality rate
 - OR/AND take blood samples for virus detection – positive result
 - **Household level:** Farm has at least one ASF positive pig confirmed





oach of investigation

CSF

ing questionnaire to collect

information

- Differential diagnosis
 - Classical swine fever (CSF)
 - Porcine reproductive and respiratory syndrome (PRRS)
 - Porcine dermatitis and nephropathy syndrome (PDNS)
 - Erysipelas
 - Aujeszky's disease
 - Salmonellosis (and other bacterial septicaemias)
 - Poisoning

Pig suffering from mycotoxin poisoning

Study size: Household/farm level

← → ↺ epitools.ausvet.com.au/output/f1a78eddfaaa1e826c874300e489415a_result.xls

Sample size to estimate a simple proportion (apparent prevalence)
Wed Dec 8 08:52:23 2021

Inputs

"Estimated Proportion" "0.01"
"Desired precision of estimate" "0.05"
"Confidence level" "0.95"
"Population size" "30000"

Results

Sample size required for specified inputs

"V1"
"Large population" 16
"Population = 30000" 16

Sample sizes for varying prevalence and precision values

"AP = 0.01"	"AP = 0.02"	"AP = 0.05"	"AP = 0.1"	"AP = 0.2"	"AP = 0.5"
"Precision = 0.01"	381	753	1825	3458	6147
"Precision = 0.02"	96	189	457	865	1537
"Precision = 0.05"	16	31	73	139	246
"Precision = 0.1"	4	8	19	35	62
"Precision = 0.2"	1	2	5	9	16

- Pigs from 3 assumed regions
- No of HH to be investigated will be calculated by Epi tool
- No of HH: **381**

Stakeholder involved

- **Investigation team:**
 - 5 trained epidemiologists
 - Veterinarians from regions
 - Village paravets
 - Head of villages
- Who provides information, who are local people to be involved
 - Farm/HH owners or farm workers
 - Pig traders
 - Farm neighbors
 - Vet drug shops
 - Abattoirs
 - Forest official/wildlife group representative
 - Sales men (meat saler/supplier) (swill?)

Source and transmission

- **ASFV source:** Wild pigs contacts (**Q4**), blood, tissues, secretions and excretions of sick and dead animals, soft ticks of the genus *Ornithodoros*
- **Farm locations:** Distance to main roads, river, markets, slaughterhouse, pig farms, infected/suspected farms, animal disposal area
- **Biosecurity practices (Q5)**
 - **Fence** surrounding the farm
 - **Net** to prevent insects (e.g. flies)
 - **Disinfection bath** at the main entrance
 - **People** enter farm daily
 - **Changing shoes or clothes** before entering farm
 - Farm keep **pets** or other animals?
 - **Boot/equipment** disinfection
 - **Disinfectants:** Kind of disinfectants/How to use? When?
 - **Workers** live inside/outside farm

Source and transmission

- **Feed and water used (Q8)**

- Source of drinking/cleaning water
- Water treatment
- Commercial/mixed feeding
- **Swill feeding?**
- Transportation feed to farm
- Distance from feed stock to pig pen
- Workers using pork in farm (bought from markets)?

- **Pig sources (Q6)**

- Introduction of piglets from a breeding farm in the last 30 days?
- Mating methods? Directly boar or insemination

- **Vaccination records (Q10)**

- Types of vaccines (CSF, FMD, Mycoplasma, PVC2, PRRS, for each type of pigs); coverage

Production loss

DESIGNATED QUESTIONNAIRE (Q11)

- Morbidity/production losses

- No of sick pig:

Body weight: _____ kg

- No of dead pig:

Body weight: _____ kg

- Cost for treatment:

- Cost for disinfection:

- Cost for culling and disposal:

- Other loss:

Estimate cost for study

No	Activity	No of people involved	Quantity	Unit (USD)	Cost per unit (USD)	Total cost (USD)
1	PILOT SURVEY (10 HH in 1 region)					1,835
	Cost for epi team					
	Travel for 2 epidemiologists to 3 regions (return-airtickets/car rental)	2	1	Lump sum	600	1,200
	Daily Allowance	2	2	Day	60	240
	Accommodation	2	2	Lump sum	50	200
	Cost for local team (3 vets/region + paravets + head villages)					
	Car/motobike rental	5	1	Lump sum	20	100
	Daily Allowance	5	1	Day	15	75
	Cost for interviewee	1	10		2	20
2	REL SURVEY (381 HH in 3 regions)					16,162
	Cost for epi team					
	Travel for 5 epidemiologists to 3 regions (return-airtickets/car rental)	5	1	Lump sum	800	4,000
	Daily Allowance	5	12	Day	60	3,600
	Accommodation	5	12	Lump sum	50	3,000
	Cost for local team (3 vets/region + paravets + head villages)					
	Car/motobike rental	5	12	Lump sum	50	3,000
	Daily Allowance	5	12	Day	30	1,800
	Cost for interviewee	1	381	Questionnaire	2	762
3	DATA ANALYSIS AND REPORT	2	5		150	1,500
	TOTAL					19,497

Summary/discussion

- Summary
 - Need to design a good questionnaire
 - Conduct pilot questionnaire to adjust if necessary
 - Identify risk factors associated with ASF in Neverland >>> recommend control measures
- Discussion
 - Sampling wild pigs? How to manage?
 - Sample size calculation: some epitool available
 - ELISA test – possible to always detect ASFv or only when the animals in fever period?

Acknowledgment

- Dr Ashish Sutar for kind facilitation and support to our group discussion/question

THANK YOU FOR YOUR ATTENTION

