



Dr Ahmad Al Majali

Professor
Faculty of Veterinary Medicine
University of Science and
Technology
Jordan
Middle East

OIE Webinar World Wildlife Day

Global wildlife health
3 March 2022

PPR in wildlife

Wildlife health globe-trotting






PPR in wildlife

- Wildlife health and diseases are **increasingly** important aspects of wildlife conservation, particularly for species that are at risk for extinction.
- The **majority of zoonotic diseases** (72%) are originating in wild animals.
- Surveillance for infectious agents in wildlife populations, and efficient investigation of wildlife disease outbreaks, are **critical to effective management of infectious diseases in wildlife**, livestock, and human populations.
- Wildlife scientists are **poorly equipped** to participate in effective disease surveillance and management in free-roaming wildlife.




PPR in wildlife

- PPR is a major **trans-boundary animal disease (TAD)** from a socio-economic point of view.
- PPR is a viral disease that affects wildlife, threatens susceptible rare **wild artiodactyl species** and is of conservation concern.



Morbillivirus


PPR isolates from wildlife are clustered in Lineage IV





PPR in wildlife

- Susceptible wildlife: **all wild ruminants**
- Clinical and pathologic findings (**similar to domesticated goats**):
 - Mucopurulent nasal discharge
 - Ulcerative keratitis along with conjunctivitis
 - Mortalities in young
 - Pinpoint greyish areas of necrosis in the buccal cavity
 - Rumen is usually congested. Abomasum exhibits tiny haemorrhagic erosions with marked congestion. Large intestine with zebra striping due to congestion and hemorrhages.
 - Consolidated lungs leading to pneumonia



Source: Munir *et al.*, 2013



(a)



(b)



(c)



(d)




(e)




(f)




(a)




(b)



(c)




(d)



(e)


Source: Prada *et al.*, 2015





PPR in wildlife - spread

- PPR virus is spread by **close contact** between infected animals.
- The virus is **shed** in secretions and excretions of infected animals.
- **Aerosol** transmission of PPR virus is also an important route of transmission.
- Animals are considered **infectious during the incubation period**, which might range from 2 to 10 days.
- No development of carriers



PPR - distribution



OIE official status (May 2018)



- OIE Members and zones recognised as free from PPR
- PPR-free status suspended

PPR GCES stages as self-assessed by the countries during Roadmap meetings in 2015-2018

- 4 ■ 3 ■ 2 ■ 1

- Countries and zones without an official PPR status and which PPR-GCES stages have not been assessed

Source: OIE, 2018




In 2018, first ever cases of PPR were reported in Burundi and Bulgaria, the first EU country affected by the disease.



PPR in wildlife - distribution

Wildlife only


2005-2019



34 countries reporting disease present




PPR in wildlife - passive surveillance





Situation in the Middle East region

Tabuk



Country	Wild animal species
UAE	<ul style="list-style-type: none">- Antelopes- Rheem gazelles,- Arabian mountain gazelles,- Springbuck, <i>Antidorcas marsupialis</i>,- Nubian ibex (<i>Capra nubiana</i>),- Barbary sheep (<i>Ammotragus lervia</i>),- Afghan Markhor goat (<i>Capra falconeri</i>)
Iran	<ul style="list-style-type: none">- Fallow Deer
Occupied Palestinian Territory	<ul style="list-style-type: none">- Fallow Deer- Mountain gazelle- Nubian ibex- Wild boar
KSA	NA

PPR in wildlife and effect on ecosystem health and biodiversity



- PPRV outbreaks in free-ranging wild artiodactyls can result in severe mortality and threaten wildlife populations and ecosystem stability
- Serological responses to PPRV in wildlife indicate widespread spillover at the wildlife-livestock interface
- The expansion of PPR into free-ranging wildlife negatively impact biodiversity and dim the vision of a PPR-free world by 2030
- Current surveillance for wildlife disease usually targets diseases that affect humans or livestock, not those impacting wildlife populations.



PPR eradication in wildlife: Global strategy

PPR Global Eradication Plan (2017-2021)

Component 1 - Promoting an enabling environment and reinforcing veterinary capacities

- 1.1: PPR strategy and technical plans
- 1.2: Stakeholder awareness and engagement
- 1.3: Legal framework
- 1.4: Strengthening veterinary services

Component 2 - Support to the diagnostic and surveillance systems

- 2.1: Epidemiological assessment
- 2.2: Strengthening surveillance systems and laboratory capacities
- 2.3: Regional epidemiology and laboratory networks

Component 3 - Measures supporting PPR eradication

- 3.1: Vaccination and other PPR prevention and control measures
- 3.2: Demonstrating PPR-free status
- 3.3: Control of other small ruminant diseases in support of PPR eradication

Component 4 - Coordination and management

- 4.1: Global level
- 4.2: Regional level
- 4.3: National level

Recommendations for wildlife integration

Component 1 – Engage wildlife and veterinary agencies in PPRV eradication at wildlife-livestock interface

- 1.1: Include wildlife in PPR GEP, regional strategies, and National Strategic Plans
- 1.2: Advocate for better integration of wildlife in PPR GEP
- 1.2: Engage wildlife agencies in planning and implementation
- 1.3/1.4: Standardize guidelines for PPR management in wildlife

Component 2 - Support wildlife diagnostic and surveillance systems

- 2.1: Increase research on epidemiological role of wildlife and determinants of susceptibility
- 2.2: Standardize guidelines for PPRV diagnostic tools in wildlife
- 2.1/2.2: Improve wildlife health surveillance, including via ecological monitoring and participatory methods
- 2.3: Include wildlife in regional epidemiology and laboratory networks

Component 3 - Integrated PPRV control efforts

- 3.1: Adapt vaccination and control strategies to the presence of susceptible and significant wildlife populations
- 3.1: Consider the entire community of susceptible host
- 3.2: Jointly monitor the effectiveness of PPRV control measures in livestock and wildlife
- 3.3: Monitor overall impact on livestock, wildlife, and ecosystem health

Component 4 - Coordination and management

- 4.1: Create a specialized group on the wildlife-livestock interface in PPR GREN
- 4.2: Incorporate wildlife in the European Food Safety Authority study on the risk for PPRV incursion in the EU
- 4.1/4.2/4.3: Ensure financial resource mobilization for the wildlife components of national, regional, and global strategies

